# CUMULATIVE SUBJECT INDEX

## Environmental Entomology, Volume 16, 1987


*Abies fraseri*, *Adelges piceae*, 712
  monoterpenes, 712
*Abutilon theophrasti*, *Niesthrea louisianica*, 963
  reproduction, 963
*Acanthoscelides obtectus*, *Phaseolus acutifolius*, 1067
  *Phaseolus vulgaris*, 1067
Acari, apples, 30
  community structure, 30
  *Phytoseiidae*, 30
  *Stigmaeidae*, 30
  species composition, 30
  *Tetranychidae*, 30
*Acer rubrum*, *Lymantria dispar*, 459
  nuclear polyhedrosis virus, 459
  *Pinus rigida*, 459
  *Populus tremuloides*, 459
  *Quercus rubra*, 459
Action sites, insecticides, 231
  *Ostrinia nubilalis*, 231, 779
*Aculus cornutus*, net $CO_2$, 660
  transpiration, 660
*Acyrthosiphon pisum*, biological control, 224
  establishment, 224
  overwintering success, 224
  predation, 224
*Adelges piceae*, *Abies fraseri*, 712
  monoterpenes, 712
Adult populations, biotic potential, 446
  *Diabrotica virgifera virgifera*, 446
  models, 446
Aerial dispersal, *Oligonychus pratensis*, 928
  *Tetranychus urticae*, 928
*Agromyza frontella*, alfalfa, 575
  life tables, 145
  population dynamics, 145
  simulation model, 575
*Agrotis ipsilon*, cotton, 743
  feeding damage, 743
  *Gossypium hirsutum*, 743
  Microsporidia, 1228
  *Vairimorpha necatrix*, 1228
*Aleurocanthus woglumi*, *Amitus hesperidum*, 140
  biological control, 140
  *Encarsia opulenta*, 140
  *Encarsia smithi*, 140
  hyperparasitism, 140
Alfalfa, Agromyza frontella, 575
  *Bruchophagus roddi*, 774
  *Hypera postica*, 195
  interaction, 1175
  potato leafhopper, 1175
  simulation model, 575
  trichomes, 195
Almonds, *Osmia cornuta*, 664
  pollination, 664
Alternate host plants, cotton pests, 1266
  *Lygus lineolaris*, 1266
*Amblyseius fallacis*, diapause induction, 72
*Amblyseius limonicus*, biological control, 1012
  cassava mites, 1012
  *Mononychellus progresivus*, 545
  natural enemies, 545
  *Oligota minuta*, 545
  permethrin, 545
  *Tetranychus urticae*, 545
*Amitus hesperidum*, *Aleurocanthus woglumi*, 140
  biological control, 140
  *Encarsia opulenta*, 140
  *Encarsia smithi*, 140
  hyperparasitism, 140
*Amyelois transitella*, *Cadra figulilella*, 630
  *Ephestia elutella*, 630
  nonoccluded virus, 630
*Anagrus mutans*, *Anagrus silwoodensis*, 825
  host effect, 825
*Anagrus silwoodensis*, *Anagrus mutans*, 825
  host effect, 825
*Anasa tristis*, biological control, 1324
  diapause, 269
  gregarious parasitoids, 1324
  reproduction, 269
  voltinism, 269
*Anastatus semiflavidus*, *Hemileuca oliviae*, 1035
  parasitoids, 1035
*Anthonomus grandis*, ovipositional behavior, 951
*Anticarsia gemmatalis*, *Nabis roseipennis*, 1330
  nuclear polyhedrosis virus, 691, 1330
  virus transmission, 691
Antixenosis, host-plant resistance, 764
  *Otiorhynchus sulcatus*, 764
*Apanteles aristoteliae*, *Argyrotaenia citrana*, 507
  benomyl, 507
*Aphelinus perpallidus*, *Monellia caryella*, 877
  reproductive response, 877
*Apis mellifera*, apples, 309
  bee distribution, 922
  *Helianthus annuus*, 922
  model, 309
  pollination, 309
Apples, Acari, 30
  *Apis mellifera*, 309
  community structure, 30
  damage, 569
  model, 309
  *Panonychus ulmi*, 569
  Phytoseiidae, 30
  *Platynota idaeusalis*, 786
  pollination, 309
  Stigmaeidae, 30
  species composition, 30
  Tetranychidae, 30
Area-wide management, cotton agroecosystem, 379
  *Lygus lineolaris*, 379
  predators, 379
  weed hosts, 379
*Argyrogramma verruca*, sex pheromones, 1030
*Argyrotaenia citrana*, *Apanteles aristoteliae*, 507
  benomyl, 507
  pheromone traps, 59
population dynamics, 59
*Artogeia rapae*, cabbages, 680
  feeding, 680
  *Trichoplusia ni*, 680
Assimilation rate, *Aculus cornutus*, 660
  net $CO_2$, 660
  transpiration, 660
Assortative mating, *Culex tarsalis*, 405
  colonization, 405
*Atherigona soccata*, oviposition stimulant, 967
  population dynamics, 910
  sorghum, 967
Atmospheric $CO_2$, *Phaseolus lunata*, 639
  *Trichoplusia ni*, 639
Attractants, *Diabrotica*, 126
  eugenol, 126
  evolution, 126
  kairomones, 126
Avian predators, *Orgyia pseudotsugata*, 90
  predation, 90

*Baccharis*, ecology, 887
  insects, 887
Bait formulations, *Heliothis virescens*, 605
  pheromone dispensers, 605
  polyvinyl chloride, 605
*Bangasternus fausti*, *Bangasternus orientalis*, 645
  biological control, 645
  *Centaurea*, 645
*Bangasternus orientalis*, *Bangasternus fausti*, 645
  biological control, 645
  *Centaurea*, 645
*Beauveria bassiana*, soil columns, 178
  vertical movement, 178
Bee distribution, *Apis mellifera*, 922
  *Helianthus annuus*, 922
Behavior, conditioning, 471
  cotton, 471
  *Gossypium hirsutum*, 471
  Haematobia irritans, 1107
  *Heliothis zea*, 471
Behavioral resistance, *Haematobia irritans*, 467
  horn flies, 467
Beneficial insects, *Coccinella septempunctata*, 368
  dispersion, 368
Benomyl, *Apanteles aristoteliae*, 507
  *Argyrotaenia citrana*, 507
Binary response, *Rhagoletis pomonella*, 319
  repeated measures, 319
  statistics, 319
Bioassay, coniine, 656
  *Heliothis zea*, 656
Biological control, *Acyrthosiphon pisum*, 224
  *Aleurocanthus woglumi*, 140
  *Amblyseius limonicus*, 1012
  *Amitus hesperidum*, 140
  *Anasa tristis*, 1324
  *Bangasternus fausti*, 645
  *Bangasternus orientalis*, 645
  *Brachymeria ovata*, 1207
  *Carduus*, 111
  cassava mites, 1012
  *Centaurea*, 645
  *Cirsium*, 111
  Cochliomyia hominivorax, 599
  *Coleophora*, 1027
  cyromazine, 828
  *Encarsia opulenta*, 140
  *Encarsia smithi*, 140
  establishment, 224
  *Fannia* sp., 769
  gregarious parasitoids, 1324
  host crops, 339
  host preference, 339
  hyperparasitism, 140
  *Leptinotarsa decemlineata*, 633
  *Liriomyza* spp., 339
  *Liriomyza trifolii*, 832
  *Musca domestica*, 769, 828
  mass rearing, 599
  nematodes, 769
  *Ophyra aenescens*, 828

*Oplomus dichrous*, 633
ornamentals, 832
overwintering success, 224
parasites, 198
parasitoids, 339, 1207
predation, 224
predators, 633
*Rhinocyllus conicus*, 111
*Rhopalomyia californica*, 198
*Salsola*
spatial density dependence, 198
thermal ecology, 599
thistle, 111
*Torymus koebelei*, 198
Bionomics, *Delia florilega*, 881
Biotic potential, adult populations, 446
*Diabrotica virgifera virgifera*, 446
models, 446
Biotypes, broken resistance, 1163
esterase, 676
polymorphism, 519
*Schizaphis graminum*, 519, 676, 1163
Blacklight traps, insect monitoring, 154
*Ostrinia nubilalis*, 154
pheromone traps, 154
water-pan traps, 154
*Blatta orientalis*, cockroach parasites, 938
distribution, 731
movement, 731
*Prosevania punctata*, 938
Bollworm, *Heliothis zea*, 1296
pheromone traps, 1296
*Brachymeria ovata*, biological control, 1207
parasitoid, 1207
Broken resistance, biotypes, 1163
*Schizaphis graminum*, 1163
*Bruchophagus roddi*, alfalfa, 774
*Bruchus pisorum*, *Eupteromalus leguminis*, 653
parasitization, 653
peas, 653
Buprestids, *Diaphnocoris chlorionis*, 996
honeylocust, 996

Cabbages, *Artogeia rapae*, 680
*Trichoplusia ni*, 680
*Cadra figulilella*, *Amyelois transitella*, 630
*Ephestia elutella*, 630
nonoccluded virus, 630
*Campoletis sonorensis*, host plant resistance, 1181
*Spodoptera frugiperda*, 1181
Cannibalism, feeding indices, 672
manipulation, 1082
*Plagiodera versicolora*, 1082
*Spodoptera frugiperda*, 672
*Carcinops pumilio*, dispersal, 415
predation, 415
*Carduus*, biological control, 111
*Cirsium*, 111
*Rhinocyllus conicus*, 111
thistle, 111
*Carya illinoensis*, gas exchange, 264
pecans, 264
*Phylloxera*, 264
photosynthesis, 264
Cassava mites, *Amblyseius limonicus*, 1012
biological control, 1012
*Centaurea*, *Bangasternus fausti*, 645
*Bangasternus orientalis*, 645
biological control, 645
Cerambycidae, development time, 563
*Metrosideros polymorpha*, 563
*Plagithmysus bilineatus*, 563
*Ceratitis capitata*, host infestations, 1273
occurrence patterns, 1273
*Ceratocystis wageneri*, *Pseudotsuga menziesii*, 1310
root beetles, 1310
Cereal aphids, distribution, 896
sampling, 896
Chemoorientation, *Diabrotica virgifera virgifera*
host location, 975
*Choristoneura fumiferana*, fecundity, 1291
oogenesis, 1291
*Choristoneura occidentalis*, cone damage, 304
defoliation, 304
fecundity, 551
oocytes, 551
Chrysanthemums, *Liriomyza trifolii*, 832
Circadian rhythms, *Heliothis virescens*, 25
larvae, 25
*Trichoplusia ni*, 25
*Cirsium*, biological control, 111
*Carduus*, 111
*Rhinocyllus conicus*, 111
thistle, 111
Climatic release, forest insect defoliator, 246
*Metrocampa guttivitta*, 246
Palmer drought index, 246
*Cnaphalocrocis medinalis*, *Goniosus triangulifer*, 1283
*Marasmia patnalis*, 1283
*Coccinella septempunctata*, beneficial insects, 368
dispersion, 368
Cochliomyia hominivorax, biological control, 599
mass rearing, 599
thermal ecology, 599
Cockroach parasites, *Blatta orientalis*, 938
*Prosevania punctata*, 938
Cohabitation, colonization, 390
*Pinus taeda*, 390
Scolytidae, 390
*Coleophora*, biological control, 1027
*Salsola*, 1027
Coleoptera, strip-mines, 168
succession patterns, 168
trophic structure, 168
Colonies, queen size, 649
*Vespula maculifrons*, 649

Colonization, assortative mating, 405
*Culex tarsalis*, 405
cohabitation, 390
*Pinus taeda*, 390
Scolytidae, 390
Color response, flight activity, 1199
Leptinotarsa decemlineata, 1199
Compensation, *Liriomyza trifolii*, 374
lima beans, 374
photosynthesis, 374
Compositae, *Homoeosoma electellum*, 586
host resistance, 586
pest management, 586
Conditioning, behavior, 471
cotton, 471
*Gossypium hirsutum*, 471
*Heliothis zea*, 471
Cone damage, *Choristoneura occidentalis*, 304
defoliation, 304
Coniine, bioassay, 656
*Heliothis zea*, 656
Control, dung beetles, 1135
*Haematobia irritans*, 1135
Corn, Elateridae, 475
rubidium, 475
*Spodoptera frugiperda*, 453
sampling, 453
Taylor power function, 453
Corn kernels, *Heliothis zea*, 535
protein content, 535
simulated damage, 535
*Corythucha cydoniae*, *Cotoneaster* spp., 365
insect resistance, 365
oviposition, 365
*Cotoneaster* spp., *Corythucha cydoniae*, 365
insect resistance, 365
oviposition, 365
Cotton, *Agrotis ipsilon*, 743
behavior, 471
conditioning, 471
croton, 355
*Gossypium hirsutum*, 471, 743
*Heliothis zea*, 420, 471
infestation foci, 614
*Oxyopes salticus*, 1114
*Peucetia*, 355
*Pseudatomoscelis*, 355
spider mites, 614
*Tetranychus* spp., 614
*Trichogramma pretiosum*, 420
Cotton agroecosystem, area-wide management, 379
*Lygus lineolaris*, 379
predators, 379
weed hosts, 379
Cotton pests, alternate host plants, 1266
*Lygus lineolaris*, 1266
*Crematogaster punctulata*, *Hemileuca oliviae*, 814
natural enemies, 814
predation, 814
Croton, cotton, 355
*Peucetia*, 355
*Pseudatomoscelis*, 355
*Cryptolestes ferrugineus*, grain moisture, 77
parasites, 77
*Rhyzopertha dominica*, 77
temperature, 77
Cucurbits, *Diaphania hyalinata*, 1062
*Diaphania nitidalis*, 1062
*Culex tarsalis*, assortative mating, 405
colonization, 405
Cultural practices, *Heliothis zea*, 237
practices, soybeans, 237
stage-specific survival, 237
Cyromazine, biological control, 828
*Musca domestica*, 828
*Ophyra aenescens*, 828

Damage, apples, 569
*Panonychus ulmi*, 569
Defoliation, *Choristoneura occidentalis*, 304
cone damage, 304
development rate, 202
*Epilachna varivestis*, 274
*Lymantria dispar*, 202
microhabitat selection, 202
pinto beans, 274
regression, 274
thermal ecology, 202
Degree-days, *Psila rosae*, 503
*Delia florilega*, bionomics, 881
*Delia platura*, host acceptance, 400
oviposition, 400
reproduction, 400
*Dendroctonus frontalis*, pines, 9
response, 9
susceptibility, 9
*Dendroctonus ponderosae*, lodgepole pine, 738
stand density index, 738
Density, *Hemileuca oliviae*, 945
Development, *Hylobius pales*, 956
*Ips calligraphus*, 484
life tables, 296
mortality, 484
*Papaipema nebris*, 296
pine regeneration pests, 956
population dynamics, 296
temperature effects, 484
Development rates, defoliation, 202
*Lymantria dispar*, 202, 556
microhabitat selection, 202
phenology, 556
temperature, 990
thermal ecology, 202
*Thyridopteryx ephemeraeformis*, 1141
*Trypoxylon*, 990
Development time, Cerambycidae, 563

*Metrosideros polymorpha*, 563
*Plagithmysus bilineatus*, 563
*Diabrotica*, attractants, 126
eugenol, 126
evolution, 126
kairomones, 126
*Diabrotica virgifera virgifera*, adult populations, 446
biotic potential, 446
chemoorientation, 975
distribution, 1193
host location, 975
models, 446
orientation, 1193
Diapause, *Anasa tristis*, 269
*Haematobia irritans*, 1092
photoperiod, 1092
reproduction, 269
voltinism, 269
Diapause induction, *Amblyseius fallacis*, 72
*Diaphania hyalinata*, cucurbits, 1062
*Diaphania nitidalis*, 1057, 1062
population dynamics, 1057
*Diaphania nitidalis*, cucurbits, 1062
*Diaphania hyalinata*, 1057, 1062
larval dispersion, 539
population dynamics, 1057
sampling, 539
*Diaphnocoris chlorionis*, buprestids, 996
honeylocust, 996
*Diaprepes abbreviatus*, *Neoaplectana carpocapsae*, 987
*Neoaplectana glaseri*, 987
Diaspididae, chlorosis, 1314
photosynthesis, 1314
Diet switching, *Heliothis zea*, 206
*Dioryctria reniculelloides*, monitoring, 905
sex pheromones, 905
Diptera, evolution, 847
Homoptera, 847
Dispersal, *Carcinops pumilio*, 415
*Lygus hesperus*, 687
mark/recapture, 211
population dynamics, 687
predation, 415
*Tabanus abactor*, 211
Dispersion, beneficial insects, 368
*Coccinella septempunctata*, 368
*Macrocheles muscaedomesticae*, 159
*Musca domestica*, 159
*Neochetina eichhorniae*, 699
waterhyacinth, 699
Distribution, *Blatta orientalis*, 731
cereal aphids, 896
*Diabrotica virgifera virgifera*, 1193
*Euonymus fortunei*, 917
Euonymus scales, 917
*Hemileuca oliviae*, 945
sampling, 896
Distributions, insecticide testing, 1
Taylor's power law, 1
Diurnal movement, leafhopper, 1103
vector, 1103
*Drosophila melanogaster*, functional response, 593
*Leptopilina boulardi*, 593
variable patch times, 593
Dung, larval and pupal survival, 722
*Musca autumnalis*, 722
Dung beetles, control, 1135
*Haematobia irritans*, 1135

Earthworms, lawn maintenance, 100
pesticides, 100
turf, 100
Eastern tent caterpillar, *Hyposoter fugitivus*, 1255
Ecological studies, *Ostrinia nubilalis*, 50
Ecology, *Baccharis*, 887
insects, 887
Economic threshold, *Empoasca fabae*, 250
*Solanum tuberosum*, 250
yield loss, 250
Edaphic conditions, fertilizer, 286
*Pulvinaria delottoi*, 286
*Pulvinariella mesembryanthemi*, 286
population dynamics, 286
*Edovum puttleri*, egg parasitoid, 1319
temperature development, 1319
Egg hatch, embryonic development, 683
*Simulium arcticum*, 683
temperature, 683
Egg parasitoid, *Edovum puttleri*, 1319
temperature development, 1319
Egg-mass dimensions, fecundity, 165
*Lymantria dispar* L., 165
regression model, 165
ELISA, peaches, 68
phony disease vectors, 68
xylem-limited bacteria, 68
Elateridae, corn, 475
rubidium, 475
Elm leaf beetle, host plant resistance, 1042
*Ulmus*, 1042
Embryonic development, egg hatch, 683
*Simulium arcticum*, 683
temperature, 683
Emergence, *Ips*, 869
reemergence, 869
*Empoasca fabae*, economic threshold, 250
*Solanum tuberosum*, 250
yield loss, 250
*Encarsia opulenta*, *Aleurocanthus woglumi*, 140
*Amitus hesperidum*, 140
biological control, 140
*Encarsia smithi*, 140
hyperparasitism, 140
*Encarsia smithi*, *Aleurocanthus woglumi*, 140
*Amitus hesperidum*, 140
biological control, 140
*Encarsia opulenta*, 140
hyperparasitism, 140
Environmental influence, *Sitophilus zeamais*, 809
*Ephestia elutella*, *Amyelois transitella*, 630
*Cadra figulilella*, 630
nonoccluded virus, 630
*Epilachna varivestis*, defoliation, 274
pinto beans, 274
regression, 274
Establishment, *Acyrthosiphon pisum*, 224
biological control, 224
overwintering success, 224
predation, 224
Esterase, biotypes, 676
*Schizaphis graminum*, 676
*Eucelatoria bryani*, fecundity, 437
*Heliothis* spp., 437
host feeding, 437
Tachinidae, 437
*Eucosma sonomana*, host selection, 1145
ponderosa pine, 1145
*Eucosma womonana*, *Helianthus*, 463
*Isophrictis similiella*, 463
pest management, 463
pheromones, 463
Eugenol, attractants, 126
*Diabrotica*, 126
evolution, 126
kairomones, 126
*Euonymus fortunei*, distribution, 917
Euonymus scales, 917
Euonymus scales, distribution, 917
*Euonymus fortunei*, 917
*Eupteromalus leguminis*, *Bruchus pisorum*, 653
parasitization, 653
pean, 653
Evolution, attractants, 126
*Diabrotica*, 126
Diptera, 847
Evolution, eugenol, 126
Homoptera, 847
kairomones, 126
selection, 15

Face flies, parasites, 902
*Fannia* sp., biological control, 769
*Musca domestica*, 769
nematodes, 769
Fecundity, *Choristoneura fumiferana*, 1291
*Choristoneura occidentalis*, 551
egg-mass dimensions, 165
*Eucelatoria bryani*, 437
*Heliothis* spp., 437
host feeding, 437
*Liriomyza trifolii*, 1001
longevity, 1001
*Lymantria dispar* L., 165
oocytes, 551
oogenesis, 1291
regression model, 165
Tachinidae, 437
Feeding, *Artogeia rapae*, 680
*Heliothis zea*, 424
nectar, 424
pigeonpea, 424
*Trichoplusia ni*, 680
Feeding behavior, host-plant resistance, 757
*Schizaphis graminum*, 757
Feeding damage, *Agrotis ipsilon*, 743
cotton, 743
*Gossypium hirsutum*, 743
Feeding indices, cannibalism, 672
*Spodoptera frugiperda*, 672
Feeding preference, host plants, 1235
*Papaipema nebris*, 1235
Fertilizer, edaphic conditions, 286
population dynamics, 286
*Pulvinaria delottoi*, 286
*Pulvinariella mesembryanthemi*, 286
Flight activity, color response, 1199
Leptinotarsa decemlineata, 1199
Foraging, *Heterotermes aureus*, 791
Foraging rates, physical environment, 802
*Solenopsis invicta*, 802
Forest insect defoliator, climatic release, 246
*Heterocampa guttivitta*, 246
Palmer drought index, 246
Functional response, *Drosophila melanogaster*, 593
host switching, 1305
*Leptopilina boulardi*, 593
spiders, 1305
variable patch times, 593

Gas exchange, *Carya illinoensis*, 264
*Phylloxera*, 264
pecans, 264
photosynthesis, 264
Genetics, host plant resistance, 327
insect biotypes, 327
pest management, 327
*Geocoris punctipes*, oviposition, 182
predator/plant relations, 182
searching behavior, 182
Geographic differences, nuclear polyhedrosis virus, 218
*Spodoptera frugiperda*, 218
*Glycine max*, *Heliothis zea*, 441
host plant suitability, 441
soybeans, 441
*Goniozus triangulifer*, *Cnaphalocrocis medinalis*, 1283
*Marasmia patnalis*, 1283
Gonotrophic development, sugar beets, 129
*Tetanops myopaeformis*, 129
*Gossypium hirsutum*, *Agrotis ipsilon*, 743
behavior, 471
conditioning, 471

cotton, 471, 743
feeding damage, 743
*Heliothis zea*, 471
Grain moisture, *Cryptolestes ferrugineus*, 77
parasites, 77
*Rhyzopertha dominica*, 77
temperature, 77
Granulosis virus, *Plodia interpunctella*, 1231
pheromones, 1231
Gregarious parasitoids, *Anasa tristis*, 1324
biological control, 1324

Habitat selection, logistic regression, 281
mosquito larvae, 281
water chemistry, 281
*Haematobia irritans*, behavior, 1107
behavioral resistance, 467
control, 1135
diapause, 1092
dung beetles, 1135
horn flies, 467
photoperiod, 1092
Height, *Neodiprion*, 1152
pheromones, 1152
*Helianthus*, *Eucosma womonana*, 463
*Isophrictis similiella*, 463
pest management, 463
pheromones, 463
*Helianthus annuus*, *Apis mellifera*, 922
bee distribution, 922
natural enemies, 1302
*Smicronyx fulvus*, 1302
*Heliothis*, bioassay, 656
overwintered density, 1076
*Heliothis* spp., *Eucelatoria bryani*, 437
fecundity, 437
host feeding, 437
Tachinidae, 437
*Heliothis virescens*, bait formulations, 605
circadian rhythms, 25
*Lygus lineolaris*, 37
larvae, 25
pheromone dispensers, 605
polyvinyl chloride, *605*
*predation*, *37*
Trichoplusia ni, *25*
Heliothis zea, behavior, 471
bollworm, 1296
conditioning, 471
coniine, 656
cotton, 420, 471
cultural practices, 237
diet switching, 206
feeding, 424
soybeans, 441
Host effect, *Anagrus mutans*, 825
*Anagrus silwoodensis*, 825
Host feeding, *Eucelatoria bryani*, 437
fecundity, 437
*Heliothis* spp., 437
Tachinidae, 437
Host infestations, *Ceratitis capitata*, 1273
occurence patterns, 1273
Host location, chemoorientation, 975
*Diabrotica virgifera virgifera*, 975
Host plant, nuclear polyhedrosis virus, 1004
*Spodoptera frugiperda*, 1004
Host plant resistance, *Campoletis sonorensis*, 1181
elm leaf beetle, 1042
genetics, 327
insect biotypes, 327
pest management, 327
*Schizaphis graminum*, 1007
*Spodoptera frugiperda*, 1181
*Triticum aestivum*, 1007
*Ulmus*, 1042
Host plant suitability, *Glycine max*, 441
*Heliothis zea*, 441
soybeans, 441
Host plants, feeding preference, 1235
*Papaipema nebris*, 1235
*Spissistilus festinus*, 513
Host preference, biological control, 339
host crops, 339
*Liriomyza* spp., 339
parasitoids, 339
Host range, lepidopterous pests, 855
*Microplitis rufiventris*, 855
Host resistance, Compositae, 586
*Homoeosoma electellum*, 586
pest management, 586
Host selection, *Eucosma sonomana*, 1145
host susceptibility, 984
*Lygus hesperus*, 984
Host susceptibility, host selection, 984
*Lygus hesperus*, 984
onions, 859
*Thrips tabaci*, 859
Host switching, functional response, 1305
spiders, 1305
Host transfer, insect races, 979
*Rhinocyllus conicus*, 979
Host-plant resistance, antixenosis, 764
feeding behavior, 757
numerical taxonomy, 839
*Otiorhynchus sulcatus*, 764
*Schizaphis graminum*, 757, 839
*Hylobius pales*, development, 956
pine regeneration pests, 956
*Hypera postica*, alfalfa, 195
*Medicago*, 708
resistance, 708
trichomes, 195
Hyperparasitism, *Aleurocanthus woglumi*, 140
*Amitus hesperidum*, 140
biological control, 140
*Encarsia opulenta*, 140
*Encarsia smithi*, 140
*Hyposoter fugitivus*, eastern tent caterpillar, 1255

Infestation foci, cotton, 614
spider mites, 614
*Tetranychus* spp., 614
Infestation prediction, *Heliothis zea*, 821
sweet corn, 821
Insect behavior, *Muscidifurax raptor*, 481
*Muscidifurax zaraptor*, 481
parasitoid, 481
Insect biotypes, genetics, 327
host plant resistance, 327
pest management, 327
Insect monitoring, blacklight traps, 154
*Ostrinia nubilalis*, 154
pheromone traps, 154
water-pan traps, 154
Insect pest management, Monte Carlo procedures, 345
sequential probability ratio test, 345
sequential sampling, 345
Insect races, host transfer, 979
*Rhinocyllus conicus*, 979
Insect resistance, *Corythucha cydoniae*, 365
*Cotoneaster* spp., 365
oviposition, 365
*Pectinophora gossypiella*, 84
sex pheromones, 84
sticky traps, 84
Insecticide testing, distributions, 1
statistics, 1
Taylor's power law, 1
Insecticides, action sites, 231
*Ostrinia nubilalis*, 231
Insects, *Baccharis*, 887
ecology, 887
Integrated control, herbicides, 864
*Trichosirocalus horridus*, 864
thistle, 864
Interaction, alfalfa, 1175
potato leafhopper, 1175
*Ips*, emergence, 869
*Ips calligraphus*, development, 484
life cycle development, 497
mortality, 484
*Pinus elliottii*, 428
phloem, 428
Scolytidae, 428
temperature effects, 484, 497
*Isophrictis similiella*, *Eucosma womonana*, 463
*Helianthus*, 463
pest management, 463
pheromones, 463

Kairomones, attractants, 126
*Diabrotica*, 126
eugenol, 126
evolution, 126

Larvae, circadian rhythms, 25
*Heliothis virescens*, 25
*Trichoplusia ni*, 25
Larval and pupal survival, dung, 722
*Musca autumnalis*, 722
Larval dispersion, *Diaphania nitidalis*, 539
sampling, 539
Lawn maintenance, earthworms, 100
pesticides, 100
turf, 100
Leafhopper, diurnal movement, 1103
Lepidopterous pests, host range, 855
*Microplitis rufiventris*, 855
*Leptinotarsa decemlineata*, biological control, 633
color response, 1199
flight activity, 1199
*Oplomus dichrous*, 633
plant diversity, 1019
population ecology, 1019
population trends, 135
predators, 633
tillage, 135
tomatoes, 135
*Leptopilina boulardi*, *Drosophila melanogaster*, 593
functional response, 593
variable patch times, 593
Life cycle development, *Ips calligraphus*, 497
temperature effects, 497
Life tables, *Agromyza frontella*, 145
development, 296
*Lygus hesperus*, 45
*Orgyia pseudotsugata*, 1217
*Papaipema nebris*, 296
*Phaseolus vulgaris*, 45
plant resistance, 45
population dynamics, 145, 296, 1217
Light traps, negative binomial, 259
*Scotinophara coarctata*, 259
sequential sampling, 259
Lima beans, compensation, 374
*Liriomyza trifolii*, 374
photosynthesis, 374
*Liriomyza* spp., biological control, 339
host crops, 339
host preference, 339
parasitoids, 339
*Liriomyza trifolii*, biological control, 832
chrysanthemums, 832
compensation, 374
fecundity, 1001
lima beans, 374
longevity, 1001
ornamentals, 832
photosynthesis, 374

Lodgepole pine, Dendroctonus ponderosae, 738
  stand density index, 738
Logistic regression, habitat selection, 281
  mosquito larvae, 281
  water chemistry, 281
Longevity, fecundity, 1001
  *Liriomyza trifolii*, 1001
Low temperature, *Myzus persicae*, 796
*Lygus hesperus*, dispersal, 687
  host selection, 984
  host susceptibility, 984
  life tables, 45
  *Phaseolus vulgaris*, 45
  plant resistance, 45
  population dynamics, 687
*Lygus lineolaris*, alternate host plants, 1266
  area wide management, 379
  cotton agroecosystem, 379
  cotton pests, 1266
  *Heliothis virescens*, 37
  predation, 37
  predators, 379
  weed hosts, 379
*Lymantria dispar*, *Acer rubrum*, 459
  defoliation, 202
  development rates, 202, 556
  microhabitat selection, 202
  nuclear polyhedrosis virus, 459
  phenology, 556
  *Pinus rigida*, 459
  *Populus tremuloides*, 459
  *Quercus rubra*, 459
  thermal ecology, 202
*Lymantria dispar* L., egg-mass dimensions, 165
  fecundity, 165
  regression model, 165

*Macrocentrus grandii*, *Nosema pyrausta*, 1071
  *Ostrinia nubilalis*, 1071
  survivorship, 1071
*Macrocheles muscaedomesticae*, dispersion, 159
  *Musca domestica*, 159
Manipulation, cannibalism, 1082
  *Plagiodera versicolora*, 1082
*Marasmia pathalis*, *Cnaphalocrocis medinalis*, 1283
  *Goniozus triangulifer*, 1283
Mark/recapture, dispersal, 211
  *Tabanus abactor*, 211
Mass rearing, biological control, 599
  Cochliomyia hominivorax, 599
  thermal ecology, 599
*Medicago*, *Hypera postica*, 708
  resistance, 708
*Melanocallis caryaefoliae*, pecan energy drain, 1045
*Melanoplus femurrubrum*, polymorphic development, 120
  voltinism, 120
*Metrosideros polymorpha*, Cerambycidae, 563
  development time, 563
  *Plagithmysus bilineatus*, 563
Microflora, *Prostephanus truncatus*, 618
  *Rhyzopertha dominica*, 618
  stored corn, 618
Microhabitat selection, defoliation, 202
  development rate, 202
  *Lymantria dispar*, 202
  thermal ecology, 202
*Microplitis rufiventris*, host range, 855
  lepidopterous pests, 855
Microsporidia, *Agrotis ipsilon*, 1228
  *Valrimorpha necatrix*, 1228
Migration, *Heliothis zea*, 1148
  *Plathypena scabra* (F.), 1169
  pollen, 1148
  trajectories, 1169
Migratory pathways, trajectories, 1244
  wind transport, 1244
Models, adult populations, 446
  *Apis mellifera*, 309
  apples, 309
  biotic potential, 446
  *Diabrotica virgifera virgifera*, 446
  pollination, 309
*Monellia caryella*, *Aphelinus perpallidus*, 877
  *Melanocallis caryaefoliae*, 1045
  *Monelliopsis pecanis*, 1045
  pecan energy drain, 1045
  reproductive response, 877
*Monelliopsis pecanis*, *Melanocallis caryaefoliae*, 1045
  *Monellia caryella*, 1045
  pecan energy drain, 1045
Monitoring, *Diorycteria reniculelloides*
  sex pheromones, 905
*Mononychellus progresivus*, *Amblyseius limonicus*, 545
  natural enemies, 545
  *Oligota minuta*, 545
  permethrin, 545
  *Tetranychus urticae*, 545
Monoterpenes, *Abies fraseri*, 712
  *Adelges piceae*, 712
Mortality, development, 484
  *Ips calligraphus*, 484
  temperature effects, 484
Mosquito larvae, habitat selection, 281
  ogistic regression, 281
  water chemistry, 281
Moth sexual communication, phenology, 1185
  pheromone traps, 1185
Movement, *Blatta orientalis*, 731
*Musca autumnalis*, dung, 722
  larval and pupal survival, 722
  sticky traps, 190
  trap design, 190
*Musca domestica*, biological control, 769, 828
  cyromazine, 828
  dispersion, 159
  *Fannia* sp., 769
  *Macrocheles muscaedomesticae*, 159
  nematodes, 769
  *Ophyra aenescens*, 828
  parasitoids, 716
  seasonal abundance, 716
*Muscidifurax raptor*, insect behavior, 481
  *Muscidifurax zaraptor*, 481
  parasitoid, 481
*Muscidifurax zaraptor*, insect behavior, 481
  *Muscidifurax raptor*, 481
  parasitoid, 481
*Myzus persicae*, low temperature, 796
  survival, 796

*Nabis roseipennis*, *Anticarsia gemmatalis*, 1330
  nuclear polyhedrosis virus, 1330
N-alkylmaleimides, pheromone perception blockage, 1130
  *Trichoplusia ni*, 1130
Natural enemies, *Amblyseius limonicus*, 545
  *Cremetogaster punctulata*, 814
  *Helianthus annuus*, 1302
  *Hemileuca oliviae*, 814
  *Mononychellus progresivus*, 545
  *Nezara viridula*, 1211
  *Oligota minuta*, 545
  permethrin, 545
  populations, 1211
  *Smicronyx fulvus*, 1302
  *Tetranychus urticae*, 545
Nectar, feeding, 424
  *Heliothis zea*, 424
  pigeonpea, 424
Negative binomial, light traps, 259
  *Scotinophara coarctata*, 259
  sequential sampling, 259
Nematodes, biological control, 769
  *Fannia* sp., 769
  *Musca domestica*, 769
*Neoaplectana carpocapsae*, *Diaprepes abbreviatus*, 987
  *Neoaplectana glaseri*, 987
*Neoaplectana glaseri*, *Diaprepes abbreviatus*, 987
  *Diaprepes abbreviatus*, 987
  *Neoaplectana carpocapsae*, 987
*Neochetina eichhorniae*, dispersion, 699
  waterhyacinth, 699
*Neodiprion*, height, 1152
  pheromones, 1152
*Nephotettix virescens*, plant age, 106
  rice, 106
  tungro virus, 106
Nesting, *Osmia cornuta*, 664
  pollination, 664
Net $CO_2$, *Aculus cornutus*, 660
  assimilation rate, 660
  transpiration, 660
*Nezara viridula*, natural enemies, 1211
  populations, 1211
*Niesthrea louisianica*, *Abutilon theophrasti*, 963
  reproduction, 963
Nonoccluded virus, *Amyelois transitella*, 630
  *Cadra figuilella*, 630
  *Ephestia elutella*, 630
*Nosema pyrausta*, horizontal transmission, 1124
  *Macrocentrus grandii*, 1071
  *Ostrinia nubilalis*, 1071, 1124
  survivorship, 1071
  sweet corn, 1124
Nuclear polyhedrosis virus, *Acer rubrum*, 459
  *Anticarsia gemmatalis*, 691, 1330
  geographic differences, 218
  host plant, 1004
  *Lymantria dispar*, 459
  *Nabis roseipennis*, 1330
  *Pinus rigida* 459
  *Populus tremuloides*, 459
  *Quercus rubra*, 459
  *Spodoptera frugiperda*, 218, 1004
  virus transmission, 691
Numerical sampling, spatial dispersion, 933
  *Urophora* sp., 933
Numerical taxonomy, host-plant resistance, 839
  *Schizaphis graminum*, 839

Occurrence patterns, *Ceratitis capitata*, 1273
  host infestation, 1273
*Oligonychus pratensis*, aerial dispersal, 928
  *Tetranychus urticae*, 928
*Oligota minuta*, *Amblyseius limonicus*, 545
  *Mononychellus progresivus*, 545
  natural enemies, 545
  permethrin, 545
  *Tetranychus urticae*, 545
Onions, host susceptibility, 859
  *Thrips tabaci*, 859
Oocytes, *Choristoneura occidentalis*, 551
  fecundity, 551
Oogenesis, *Choristoneura fumiferana*, 1291
  fecundity, 1291
*Ophyra aenescens*, biological control, 828
  cyromazine, 828
  *Musca domestica*, 828
*Oplomus dichrous*, biological control, 633
  *Leptinotarsa decemlineata*, 633
  predators, 633
*Orgyia pseudotsugata*, avian predators, 90
  life tables, 1217
  population dynamics, 1217
  predation, 90
Orientation, *Diabrotica virgifera virgifera*, 1193
Ornamentals, biological control, 832
  *Liriomyza trifolii*, 832
*Osmia*, *Pinus* pollen, 668
  pollen foraging, 668
*Osmia cornuta*, almonds, 664

nesting, 664
pollination, 664
*Ostrinia nubilalis*, action sites, 231, 779
blacklight traps, 154
ecological studies, 50
horizontal transmission, 1124
insect monitoring, 154
insecticides, 231, 779
*Macrocentrus grandii*, 1071
*Nosema pyrausta*, 1071, 1124
pheromone traps, 154
spiders, 779
survivorship, 1071
sweet corn, 1124
water-pan traps, 154
*Otiorhynchus sulcatus*, antixenosis, 764
host-plant resistance, 764
Overwintered density, *Heliothis*, 1076
Overwintering success, *Acyrthosiphon pisum*, 224
biological control, 224
establishment, 224
predation, 224
Oviposition, *Corythucha cydoniae*, 365
*Cotoneaster* spp., 365
*Delia platura*, 400
*Geocoris punctipes*, 182
host acceptance, 400
insect resistance, 365
predator/plant relations, 182
reproduction, 400
searching behavior, 182
*Sitophilus zeamais*, 809
Oviposition site selection, *Heliothis zea*, 625
soybeans, 625
Oviposition stimulant, *Atherigona soccata*, 967
sorghum, 967
Ovipositional behavior, *Anthonomus grandis* 951
*Oxyopes salticus*, cotton, 1114
prey capture rate, 1114

Palmer drought index, climatic release, 246
forest insect defoliator, 246
*Heterocampa guttivitta*, 246
*Panonychus ulmi*, apples, 569
damage, 569
*Papaipema nebris*, development, 296
feeding preference, 1235
host plants, 1235
life tables, 296
population dynamics, 296
Parasites, biological control, 198
*Cryptolestes ferrugineus*, 77
face flies, 902
grain moisture, 77
*Rhopalomyia californica*, 198
*Rhyzopertha dominica*, 77
spatial density dependence, 198
*Torymus koebelei*, 198
temperature, 77
Parasitization, *Bruchus pisorum*, 653
*Eupteromalus leguminis*, 653
peas, 653
sugarcane, 360
Trichogrammatidae, 360
Parasitoids, *Anasstatus semiflavidus*, 1035
biological control, 339, 1207
*Brachymeria ovata*, 1207
*Hemileuca oliviae*, 1035
host crops, 339
host preference, 339
insect behavior, 481
*Liriomyza* spp., 339
*Musca domestica*, 716
*Muscidifurax raptor*, 481
*Muscidifurax zaraptor*, 481
seasonal abundance, 716
Peaches, ELISA, 68
phony disease vectors, 68
xylem-limited bacteria, 68
Peas, *Bruchus pisorum*, 653
*Eupteromalus leguminis*, 653
parasitization, 653
Pecan energy drain, *Melanocallis caryaefoliae*, 1045
*Monellia caryella*, 1045
*Monelliopsis pecanis*, 1045
Pecans, *Carya illinoensis*, 264
gas exchange, 264
*Phylloxera*, 264
photosynthesis, 264
*Pectinophora gossypiella*, insecticide resistance, 84
sex pheromones, 84
sticky traps, 84
Permethrin, *Amblyseius limonicus*, 545
*Mononychellus progresivus*, 545
natural enemies, 545
*Oligota minuta*, 545
*Tetranychus urticae*, 545
Pest insects, *Phidippus audax*, 1098
predaceous insects, 1098
Pest management, Compositae, 586
*Eucosma womonana*, 463
genetics, 327
*Helianthus*, 463
*Homoeosoma electellum*, 586
host plant resistance, 327, 586
insect biotypes, 327
*Isophrictis similiella*, 463
pheromones, 463
Pesticides, earthworms, 100
lawn maintenance, 100
turf, 100
*Peucetia*, cotton, 355
*Pseudatomoscelis*, 355
*Phaseolus acutifolius*, *Acanthoscelides obtectus*, 1067
*Phaseolus vulgaris*, 1067
*Phaseolus lunata*, atmospheric $CO_2$, 639
*Trichoplusia ni*, 639
*Phaseolus vulgaris*, *Acanthoscelides obtectus*, 1067
*Lygus hesperus*, 45
life tables, 45
*Phaseolus acutifolius*, 1067
plant resistance, 45
Phenology, development rates, 556
*Lymantria dispar*, 556
moth sexual communication, 1105
pheromone traps, 1105
Pheromone dispensers, bait formulations, 605
*Heliothis virescens*, 605
polyvinyl chloride, 605
Pheromone perception blockage, N-alkylmaleimides, 1130
*Trichoplusia ni*, 1130
Pheromone trapping, *Heliothis zea*, 1241
*Rhyzopertha dominica*, 747
*Spodoptera frugiperda*, 1241
Pheromone traps, *Argyrotaenia citrana*, 59
blacklight traps, 154
bollworm, 1296
*Heliothis zea*, 1296
insect monitoring, 154
moth sexual communication, 1105
*Ostrinia nubilalis*, 154
phenology, 1105
population dynamics, 59
water-pan traps, 154
Pheromones, *Eucosma womonana*, 463
granulosis virus, 1231
height, 1152
*Helianthus*, 463
*Isophrictis similiella*, 463
*Neodiprion*, 1152
pest management, 463
*Plodia interpunctella*, 1231
*Phidippus audax*, pest insects, 1098
predaceous insects, 1098
Phloem, *Ips calligraphus*, 428
*Pinus elliottii*, 428
Scolytidae, 428
*Pholetesor ornigis*, *Pholetesor pedias*, 1157
*Phyllonorycter blancardella*, 1157
*Pholetesor pedias*, *Pholetesor ornigis*, 1157
*Phyllonorycter blancardella*, 1157
Phony disease vectors, ELISA, 68
peaches, 68
xylem-limited bacteria, 68
Photoperiod, diapause, 1092
*Haematobia irritans*, 1092
Photosynthesis, *Carya illinoensis*, 264
chlorosis, 1314
compensation, 374
Diaspididae, 1314
gas exchange, 264
*Liriomyza trifolii*, 374
lima beans, 374
pecans, 264
*Phylloxera*, 264
*Phyllonorycter blancardella*, *Pholetesor ornigis*, 1157
*Pholetesor pedias*, 1157
*Phylloxera*, *Carya illinoensis*, 264
gas exchange, 264
pecans, 264
photosynthesis, 264
Physical environment, foraging rates, 802
*Solenopsis invicta*, 802
Phytoseiidae, Acari, 30
apples, 30
community structure, 30
Stigmaeidae, 30
Phytoseiidae, species composition, 30
Tetranychidae, 30
Pigeonpea, feeding, 424
*Heliothis zea*, 424
nectar, 424
Pine regeneration pests, development, 956
*Hylobius pales*, 956
Pines, *Dendroctonus frontalis*, 9
response, 9
susceptibility, 9
Pinto beans, defoliation, 274
*Epilachna varivestis*, 274
regression, 274
*Pinus elliottii*, *Ips calligraphus*, 428
phloem, 428
Scolytidae, 428
*Pinus pollen*, *Osmia*, 668
pollen foraging, 668
*Pinus rigida*, *Acer rubrum*, 459
*Lymantria dispar*, 459
nuclear polyhedrosis virus, 459
*Populus tremuloides*, 459
*Quercus rubra*, 459
*Pinus taeda*, cohabitation, 390
colonization, 390
Scolytidae, 390
*Plagiodera versicolora*, cannibalism, 1082
manipulation, 1082
*Plagithmysus bilineatus*, Cerambycidae, 563
development time, 563
*Metrosideros polymorpha*, 563
Plant age, *Nephotettix virescens*, 106
rice, 106
tungro virus, 106
Plant diversity, *Leptinotarsa decemlineata*, 1019
population ecology, 1019
Plant resistance, *Lygus hesperus*, 45
life tables, 45
*Phaseolus vulgaris*, 45
*Plathypena scabra* (F.), migration, 1169
trajectories, 1169
*Platynota idaeusalis*, apples, 786
spatial patterns, 786
*Plodia interpunctella*, granulosis virus, 1231
pheromones, 1231

Pollen, *Heliothis zea*, 1148
  migration, 1148
Pollen foraging, *Osmia*, 668
  *Pinus* pollen, 668
Pollination, *Apis mellifera*, 309
  almonds, 664
  apples, 309
  model, 309
  *Osmia cornuta*, 664
Polymorphic development, *Melanoplus femurrubrum*, 120
  voltinism, 120
Polymorphism, biotypes, 519
  *Schizaphis graminum*, 519
Polyvinyl chloride, bait formulations, 605
  *Heliothis virescens*, 605
  pheromone dispensers, 605
Ponderosa pine, *Eucosma sonomana*, 1145
Population dynamics, *Agromyza frontella*, 145
  *Argyrotaenia citrana*, 59
Population dynamics, *Atherigona soccata*, 910
  *Diaphania hyalinata*, 1057
  *Diaphania nitidalis*, 1057
  development, 296
  dispersal, 687
  edaphic conditions, 286
  fertilizer, 286
  *Lygus hesperus*, 687
  life tables, 145, 296, 1217
  *Orgyia pseudotsugata*, 1217
  *Papaipema nebris*, 296
  *Pulvinaria delottoi*, 286
  *Pulvinariella mesembryanthemi*, 286
  pheromone traps, 59
Population ecology, *Leptinotarsa decemlineata*, 1019
  plant diversity, 1019
Population trends, *Leptinotarsa decemlineata*, 135
  tillage, 135
  tomatoes, 135
Populations, *Nezara viridula*, 1211
  natural enemies, 1211
*Populus tremuloides*, *Acer rubrum*, 459
  *Lymantria dispar*, 459
  nuclear polyhedrosis virus, 459
  *Pinus rigida*, 459
  *Quercus rubra*, 459
Potato leafhopper, alfalfa, 1175
  interaction, 1175
Predaceous insects, *Rhidippus audax*, 1098
  pest insects, 1098
Predation, *Acyrthosiphon pisum*, 224
  avian predators, 90
  biological control, 224
  *Carcinops pumilio*, 415
  *Crematogaster punctulata*, 814
  dispersal, 415
  establishment, 224
  *Heliothis virescens*, 37
  *Hemileuca oliviae*, 814
  *Lygus lineolaris*, 37
  *Orgyia pseudotsugata*, 90
  overwintering success, 224
Predator/plant relations, *Geocoris punctipes*, 182
  oviposition, 182
  searching behavior, 182
Predators, area-wide management, 379
  biological control, 633
  cotton agroecosystem, 379
  *Leptinotarsa decemlineata*, 633
  *Lygus lineolaris*, 379
  *Oplomus dichrous*, 633
  weed hosts, 379
Prepupa behavior, *Heliothis zea*, 94
  pupa survival, 94
  pupation sites, 94
Prey capture rate, *Oxyopes salticus*, 1114
*Prosevania punctata*, *Blatta orientalis*, 938
  cockroach parasites, 938
*Prostephanus truncatus*, microflora, 618
  *Rhyzopertha dominica*, 618
  stored corn, 618
Protein content, corn kernels, 535
  *Heliothis zea*, 535
  simulated damage, 535
*Pseudatomoscelis*, cotton, 355
  croton, 355
  *Peucetia*, 355
*Pseudotsuga menziesii*, root beetles, 1310
  *Ceratocystis wageneri*, 1310
*Psila rosae*, degree-days, 503
*Pulvinaria delottoi*, edaphic conditions, 286
  fertilizer, 286
  population dynamics, 286
  *Pulvinariella mesembryanthemi*, 286
*Pulvinariella mesembryanthemi*, edaphic conditions, 286
  fertilizer, 286
  *Pulvinaria delottoi*, 286
  population dynamics, 286
Pupa survival, *Heliothis zea*, 94
  prepupa behavior, 94
  pupation sites, 94
Pupation sites, *Heliothis zea*, 94
  prepupa behavior, 94
  pupa survival, 94

Queen size, colonies, 649
  *Vespula maculifrons*, 649
*Quercus rubra*, *Acer rubrum*, 459
  *Lymantria dispar*, 459
  nuclear polyhedrosis virus, 459
  *Pinus rigida*, 459
  *Populus tremuloides*, 459

Reemergence, Emergence, 869
  *Ips*, 869
Regression, defoliation, 274
  *Epilachna varivestis*, 274
  pinto beans, 274
Regression model, egg-mass dimensions, 165
  fecundity, 165
  *Lymantria dispar* L., 165
Repeated measures, binary response, 319
  *Rhagoletis pomonella*, 319
  statistics, 319
Reproduction, *Abutilon theophrasti*, 963
  *Anasa tristis*, 269
  *Delia platura*, 400
  diapause, 269
  host acceptance, 400
  *Diesthrea louisianica*, 963
  oviposition, 400
  voltinism, 269
Reproductive response, *Aphelinus perpallidus*, 877
  *Monellia caryella*, 877
Resistance, *Hypera postica*, 708
  *Medicago*, 708
Response, *Dendroctonus frontalis*, 9
  pines, 9
*Rhagoletis pomonella*, binary response, 319
  repeated measures, 319
  statistics, 319
*Rhinocyllus conicus*, biological control, 111
  *Carduus*, 111
  *Cirsium*, 111
  host transfer, 979
  insect races, 979
  temperature influence, 971
  thistle, 111
*Rhopalomyia californica*, biological control, 198
  parasites, 198
  spatial density dependence, 198
  *Torymus koebelei*, 198
*Rhyacionia zozana*, sex pheromone identification, 1287
*Rhyzopertha dominica*, *Cryptolestes ferrugineus*, 77
  grain moisture, 77
  microflora, 618
  parasites, 77
  pheromone trapping, 747
  *Prostephanus truncatus*, 618
  stored corn, 618
  temperature, 77
Rice, *Nephotettix virescens*, 106
  plant age, 106
  *Spodoptera frugiperda*, 116
  tungro virus, 106
Root beetles, *Ceratocystis wageneri*, 1310
  *Pseudotsuga menziesii*, 1310
Rubidium, corn, 475
  Elateridae, 475

*Saldula usingeri*, saline springs, 1087
  thermal springs, 1087
Saline springs, *Saldula usingeri*, 1087
  thermal springs, 1087
*Salsola*, biological control, 1027
  *Coleophora*, 1027
Sampling, cereal aphids, 896
  corn, 453
  *Diaphania nitidalis*, 539
  distribution, 896
  larval dispersion, 539
  *Spodoptera frugiperda*, 453
  summer squash, 539
  Taylor power function, 453
  within/between plant, 539
*Schizaphis graminum*, biotypes, 519, 676, 1163
  broken resistance, 1163
  esterase, 676
  feeding behavior, 757
  host plant resistance, 757, 839, 1007
  numerical taxonomy, 839
  polymorphism, 519
  *Triticum aestivum*, 1007
Scolytidae, cohabitation, 390
  colonization, 390
  *Ips calligraphus*, 428
  *Pinus elliottii*, 428
  *Pinus taeda*, 390
  phloem, 428
*Scotinophara coarctata*, light traps, 259
  negative binomial, 259
  sequential sampling, 259
Searching behavior, *Geocoris punctipes*, 182
  oviposition, 182
  predator/plant relations, 182
Selection, evolution, 15
Sequential probability ratio test, insect pest management, 345
  Monte Carlo procedures, 345
  sequential sampling, 345
Sequential sampling, insect pest management, 345
  light traps, 259
  Monte Carlo procedures, 345
  negative binomial, 259
  *Scotinophara coarctata*, 259
  sequential probability ratio test, 345
Sex pheromone identification, *Rhyacionia zozana*, 1287
Sex pheromones, *Argyrogramma verruca*, 1030
  *Diarsetria reniculelloides*, 905
  insecticide resistance, 84
  monitoring, 905
  *Pectinophora gossypiella*, 84
  sticky traps, 84
Simulated damage, corn kernels, 535
  *Heliothis zea*, 535
  protein content, 535
Simulation model, Agromyza frontella, 575
  alfalfa, 575
*Simulium arcticum*, egg hatch, 683
  embryonic development, 683
  temperature, 683
*Sitophilus zeamais*, environmental influence, 809

oviposition, 809
spatial distribution, 809
*Smicronyx fulvus*, *Helianthus annuus*, 1302
natural enemies, 1302
Soil columns, *Beauveria bassiana*, 178
vertical movement, 178
Soil insecticides, soybeans, 524
tillage systems, 524
*Solanum tuberosum*, *Empoasca fabae*, 250
economic threshold, 250
yield loss, 250
*Solenopsis invicta*, foraging rates, 802
physical environment, 802
Sorghum, *Atherigona soccata*, 967
oviposition stimulant, 967
Soybeans, cultural practices, 237
*Glycine max*, 441
*Heliothis zea*, 237, 441, 625
host plant suitability, 441
oviposition site selection, 625
soil insecticides, 524
stage-specific survival, 237
tillage systems, 524
Spatial density dependence, biological control, 198
parasites, 198
*Rhopalomyia californica*, 198
*Torymus koebelei*, 198
Spatial dispersion, numerical sampling, 933
*Urophora* sp., 933
Spatial distribution, *Sitophilus zeamais*, 809
Spatial patterns, *Platynota idaeusalis*, 786
Species composition, Acari, 30
apples, 30
Phytoseiidae, 30
Stigmaeidae, 30
Tetranychidae, 30
*Sphenophorus parvulus*, survival, 752
wheatgrass, 752
Spider mites, cotton, 614
infestation foci, 614
*Tetranychus* spp., 614
Spiders, functional response, 1305
host switching, 1305
*Spissistilus festinus*, host plants, 513
*Spodoptera frugiperda*, *Campoletis sonorensis*, 1181
cannibalism, 672
corn, 453
feeding indices, 672
geographic differences, 218
*Heliothis zea*, 1241
host plant, 1004
host plant resistance, 1181
nuclear polyhedrosis virus, 218, 1004
pheromone trapping, 1241
rice, 116
sampling, 453
Taylor power function, 453
Stage-specific survival, cultural practices, 237
*Heliothis zea*, 237
soybeans, 237
Stand density index, Dendroctonus ponderosae, 738
lodgepole pine, 738
Statistics, binary response, 319
insecticide testing, 1
repeated measures, 319
*Rhagoletis pomonella*, 319
Taylor's power law, 1
Sticky traps, insecticide resistance, 84
*Musca autumnalis*, 190
*Pectinophora gossypiella*, 84
sex pheromones, 84
trap design, 190
Stigmaeidae, Acari, 30
apples, 30
community structure, 30
Phytoseiidae, 30
species composition, 30
Tetranychidae, 30
Stored corn, microflora, 618
*Prostephanus truncatus*, 618
*Rhyzopertha dominica*, 618
Strip-mines, Coleoptera, 168
succession patterns, 168
trophic structure, 168
Succession patterns, Coleoptera, 168
strip-mines, 168
trophic structure, 168
Sugar beets, gonotrophic development, 129
*Tetanops myopaeformis*, 129
Sugarcane, parasitization, 360
Trichogrammatidae, 360
Survival, *Myzus persicae*, 796
*Sphenophorus parvulus*, 752
Survivorship, *Macrocentrus grandii*, 1071
*Nosema pyrausta*, 1071
*Ostrinia nubilalis*, 1071
Susceptibility, *Dendroctonus frontalis*, 9
pines, 9
Sweet corn, *Heliothis zea*, 821
horizontal transmission, 1124
infestation prediction, 821
*Nosema pyrausta*, 1124
*Ostrinia nubilalis*, 1124
Systematic foraging, *Heterotermes aureus*, 791

*Tabanus abactor*, dispersal, 211
mark/recapture, 211
Tachinidae, *Eucelatoria bryani*, 437
fecundity, 437
*Heliothis* spp., 437
host feeding, 437
Taylor power function, corn, 453
*Spodoptera frugiperda*, 453
sampling, 453
Taylor's power law, distributions, 1
insecticide testing, 1
statistics, 1
Temperatures, *Cryptolestes ferrugineus*, 77
development rate, 990
egg hatch, 683
embryonic development, 683
grain moisture, 77
parasites, 77
*Rhyzopertha dominica*, 77
*Simulium arcticum*, 683
*Thyridopteryx ephemeraeformis*, 1141
*Trypoxylon*, 990
Temperature development, *Edovum puttleri*, 1319
egg parasitoid, 1319
Temperature effects, development, 484
*Ips calligraphus*, 484, 497
life cycle development, 497
mortality, 484
Temperature influence, *Rhinocyllus conicus*, 971
*Tetanops myopaeformis*, gonotrophic development, 129
sugar beets, 129
Tetranychidae, Acari, 30
apples, 30
community structure, 30
Phytoseiidae, 30
species composition, 30
Stigmaeidae, 30
*Tetranychus* spp., cotton, 614
infestation foci, 614
spider mites, 614
*Tetranychus urticae*, aerial dispersal, 928
*Amblyseius limonicus*, 545
*Mononychellus progresivus*, 545
natural enemies, 545
*Oligonychus pratensis*, 928
*Oligota minuta*, 545
permethrin, 545
Thermal ecology, biological control, 599
Cochliomyia hominivorax, 599
defoliation, 202
development rate, 202
*Lymantria dispar*, 202
mass rearing, 599
microhabitat selection, 202
Thermal springs, *Saldula usingeri*, 1087
saline springs, 1087
Thistle, biological control, 111
*Carduus*, 111
*Cirsium*, 111
herbicides (2,4-D), 864
integrated control, 864
*Rhinocyllus conicus*, 111
*Trichosirocalus horridus*, 864
*Thrips tabaci*, host susceptibility, 859
onions, 859
*Thyridopteryx ephemeraeformis*, development rate, 1141
temperatures, 1141
Tillage, *Leptinotarsa decemlineata*, 135
population trends, 135
tomatoes, 135
Tillage systems, soil insecticides, 524
soybeans, 524
Tomatoes, *Leptinotarsa decemlineata*, 135
population trends, 135
tillage, 135
*Torymus koebelei*, biological control, 198
parasites, 198
*Rhopalomyia californica*, 198
spatial density dependence, 198
Trajectories, migration, 1169
migratory pathways, 1244
*Plathypena scabra* (F.), 1169
wind transport, 1244
Transpiration, *Aculus cornutus*, 660
assimilation rate, 660
net $CO_2$, 660
Trap design, *Musca autumnalis*, 190
sticky traps, 190
*Trichogramma pretiosum*, cotton, 420
*Heliothis zea*, 420
Trichogrammatidae, parasitization, 360
sugarcane, 360
Trichomes, alfalfa, 195
*Hypera postica*, 195
*Trichoplusia ni*, *Artogeia rapae*, 680
atmospheric $CO_2$, 639
cabbages, 680
circadian rhythms, 25
feeding, 680
*Heliothis virescens*, 25
larvae, 25
N-alkylmaleimides, 1130
*Phaseolus lunata*, 639
pheromone perception blockage, 1130
*Trichosirocalus horridus*, herbicides (2,4-D), 864
integrated control, 864
thistle, 864
*Triticum aestivum*, host plant resistance, 1007
*Schizaphis graminum*, 1007
Trophic structure, Coleoptera, 168
strip-mines, 168
succession patterns, 168
*Trypoxylon*, development rate, 990
temperature, 990
Tungro virus, *Nephotettix virescens*, 106
plant age, 106
rice, 106
Turf, earthworms, 100
lawn maintenance, 100
pesticides, 100

*Ulmus*, elm leaf beetle, 1042
host plant resistance, 1042
*Urophora* sp., numerical sampling, 933
spatial dispersion, 933

*Vairimorpha necatrix*, Agrotis ipsilon, 1228
Microsporidia, 1228
Variable patch times, *Drosophila melanogaster*, 593
functional response, 593
*Leptopilina boulardi*, 593
Vector, diurnal movement, 1103
leafhopper, 1103
Vertical movement, *Beauveria bassiana*, 178
soil columns, 178
*Vespula maculifrons*, colonies, 649
queen size, 649
Virus transmission, *Anticarsia gemmatalis*, 691
nuclear polyhedrosis virus, 691
Voltinism, *Anasa tristis*, 269
diapause, 269
*Melanoplus femurrubrum*, 120
polymorphic development, 120
reproduction, 269

Water chemistry, habitat selection, 281
logistic regression, 281
mosquito larvae, 281
Waterhyacinth, dispersion, 699
*Neochetina eichhorniae*, 699
Water-pan traps, blacklight traps, 154
insect monitoring, 154
*Ostrinia nubilalis*, 154
pheromone traps, 154
Weed hosts, area-wide management, 379
cotton agroecosystem, 379
*Lygus lineolaris*, 379
predators, 379
Wheatgrass, *Sphenophorus parvulus*, 752
Wind transport, migratory pathways, 1244
trajectories, 1244

Xylem-limited bacteria, ELISA, 68
peaches, 68
phony disease vectors, 68

Yield loss, *Empoasca fabae*, 250
economic threshold, 250
*Solanum tuberosum*, 250

# CUMULATIVE AUTHOR INDEX

**Environmental Entomology,**
**Volume 16, 1987**


Abdel-Fattah, M., 1004
Adler, P.H., 424
Allen, D.C., 246
Alvarado-Rodriguez, B., 45
Andersen, P.C., 264, 660
Arnold, J.A., 738
Annis, B., 653
Apperson, C.S., 281
Arnold, T.B., 100
Arthur, F.H., 712
Asensio, E., 664
Asman, S.M., 405

Bailey, W.C., 296
Baker, T.C., 84
Banks, D.J., 676
Barney, R.J., 809
Bauman, T.R., 990
Bay, D.E., 159
Beers, E.H., 569
Bellinger, R.G., 120
Bellotti, A.C., 545, 1012
Benedict, J.H., 420, 471
Bentley, M.D., 967
Bergman, M.K., 975
Bloomfield, P., 281
Bolvin, G., 503
Boldt, P.E., 887
Bradley, J.R. Jr., 94, 237, 441, 625
Braun, A.R., 545, 1012
Brewer, M.J., 539
Broce, A.B., 722
Browning, H.W., 360
Buckley, P.M., 532
Bueno, R. Jr., 877
Burkholder, W.E., 747
Byford, R.L., 467

Campbell, J.B., 190, 902
Capinera, J.L., 154, 274, 859, 1019
Carey, J.R., 814
Casagrande, R.A., 556, 633
Center, T.D., 699
Chamberlain, J.D., 764
Chang, C.J., 68
Chapman, P.L., 274
Cheshire, J.M. Jr., 475
Christie, G.D., 832
Clancy, K.M., 551
Cleveland, T.C., 37
Cockfield, S.D., 917
Coffelt, M.A., 365
Cook, S.P., 9, 390, 869
Cooksey, L.M., 211
Cornell, J.A., 140, 428
Coulson, R.N., 390, 484, 497, 869
Croft, B.A., 30, 59, 507
Cross, E.A. Jr., 990
Cushing, N., 30
Cyr, T.L., 769

Dahlem, G.A., 847
Danielson, S.D., 195, 708
Davis, F.M., 532
Dean, D.A., 355
DeGrandi-Hoffman, G., 309
Demianyk, C.J., 618
DePonti, O.M.B., 327
Dgetluck, N., 319
Diehl, S.R., 319
Dorschner, K.W., 757, 1007
Doss, R.P., 764
Downes, W.L. Jr., 847
Drolet, J., 145
Drummond, F.A., 633
Duffey, S.S., 45
Dumas, B.A., 513
Dysart, R.J., 368

Eckenrode, C.J., 881
Edwards, G.B., 779
Ehler, L.E., 198
Eikenbary, R.D., 757, 1007
Eisenbach, J., 519
Elkinton, J.S., 202, 415
Elliott, N.C., 896
Epsky, N.D., 274

Fargo, W.S., 484, 497, 839
Fernandes, G.W., 15
Ferrer, E.R., 259
Fick, G.W., 575
Flamm, R.O., 390, 484, 497, 869
Flanders, R.V., 716
Fleischer, S.J., 379
Foltz, J.L., 428

Foster, K.W., 45
Foster, M.A., 743
Fowler, G.W., 345
Frank, C.J., 304
Franke, D.E., 467
Frankie, G.W., 286
Frater, A.P., 945
Fritz, G.N., 945
Fuxa, J.R., 218, 453, 1004

Gahukar, R.T., 910
Gardner, W.A., 178
Garth, J.K.L., 764
Gaylor, M.J., 379, 743
Geden, C.J., 415
Gershenzon, J., 586
Girma, M., 757
Goeden, R.D., 979, 1027
Goodenough, J.L., 605
Gould, F., 94, 327
Grace, J.K., 286
Granovsky, T.A., 747
Grant, G.G., 905
Groden, E., 633
Grodowitz, M.J., 722
Guerrero, J.M., 545, 1012
Guppy, J.C., 145
Guthrie, W.D., 50

Haack, R.A., 428
Hagstrum, D.W., 77
Hain, F.P., 9, 712
Hammond, R.B., 524
Hansen, J.D., 752
Hara, A.H., 339
Harbers, L.H., 722
Harcourt, D.G., 145
Harper, P., 129
Harrison, W., 855
Hassanali, A., 967
Havukkala, I.J., 41
Haynes, K.F., 84
Heath, R.R., 1030
Hein, G.L., 446
Heinrichs, E.A., 106
Hendricks, D.E., 605
Herms, D.A., 996
Hartlein, M.B., 593
Hoffmann, D.F., 630
Hogge, L.R., 463
Hoopingarner, R., 309
Horton, D.R., 274, 1019
Hoy, C.W., 680
Huddleston, E.W., 814, 945
Hull, L.A., 569, 786

Inayatullah, C., 839

Jarvis, J.L., 50
Jenkins, M.J., 304, 738
Johnson, G.D., 190
Johnson, K.B., 250
Johnson, M.W., 25, 339
Johnson, R.C., 757, 1007
Jones, C.G., 165
Jones, S.C., 791
Jones, V.P., 25, 832

Keaster, A.J., 475
Keating, S.T., 459
Kellen, W.R., 630
Kendall, D.M., 859
Kennedy, G.G., 327
Kieckhefer, R.W., 896
Kim, T.H., 881
King, G.G.S., 905
Kirkbride, D.M., 551
Kissam, J.B., 828
Knight, A.L., 59
Knop, N.F., 405
Koirtyohann, S.R., 475
Kok, L.T., 864, 971
Kreitner, G.L., 586
Krysan, J.L., 126

Lack, J.R., 691
Ladd, T.L. Jr., 126
Lance, D.R., 202
Landis, D.A., 94
Landolt, P.J., 1030
Lasack, P.M., 296
Legg, D.E., 809
Leigh, T.F., 45, 614
Leos-Martinez, J., 747
Li, W.-G., 84

Linduska, J.J., 135
Lockwood, J.A., 467
Logan, P.A., 556
Lopez, J.D., 420
Lord,W.D., 649
Lund, R.E., 933
Lynch, A.M., 345

Mabry, T.J., 586
MacMahon, J.A., 168
Maddox, D.M., 645
Manglitz, G.R., 195, 708
Margolies, D.C., 928
Martens, B., 374
Martinat, P.J., 246
Mason, R.R., 90
Mayo, Z.B., 676
McArdle, T.M., 390
McClure, M.S., 224
McCutcheon, G.S., 855
McGovern, W.L., 951
McLeod, P., 796
McNeil, J.N., 145
Meagher, R.L. Jr., 786
Melton, C.W., 360
Merchant, M.E., 716
Meyer, J.A., 769
Milby, M.M., 405
Miller, J.R., 41, 400
Miller, T.A., 84
Mitchell, E.B., 951
Mitchell, F.L., 453
Mittler, T.E., 519
Mizell, R.F. III, 264, 660
Moffett, J.O., 774
Moore, K.E.B., 165
Moratorio, M.S., 825
Morrison, R.K., 420
Mueller, A.J., 513
Mullens, B.A., 769
Muller, H., 1027

Naranjo, S.E., 182
Nechols, J.R., 269
Nettles, W.C. Jr., 437
Nielsen, D.G., 996
Nitao, J.K., 656
Nolan, M.P. III, 828
Nowierski, R.M., 933
Nutting, W.L., 791
Nyffeler, M., 355

Ohno, N., 586
O'Keefe, L.E., 653
Osbrink, W.L.A., 639
Owens, J.C., 814, 945

Pantoja, A., 116
Parmenter, R.R., 168
Parrella, M.P., 832
Pedigo, L.P., 296
Pemberton, R.W., 111
Pickett, C.H., 614
Pienkowski, R.L., 120
Pilcher, S.D., 154
Porter, S.D., 802
Potter, D.A., 100, 917
Prevost, Y.H., 905
Price, P.W., 15
Prokopy, R.J., 319
Pulcer, R., 309
Pulley, P.E., 390, 484, 869

Radcliffe, E.B., 250
Raffa, K.F., 672
Rapusas, H.R., 106
Reisen, W.K., 405
Richman, D.B., 814
Richter, A.R., 1004
Ricker, D.W., 1027
Robbins, T.O., 887
Robinson, J.F., 116
Robinson, W.H., 731, 938
Rodriguez, J.G., 809
Rogers, C.E., 463, 586
Rosenthal, S.S., 111
Rosset, P.M., 481
Roth, R.R., 649
Rubink, W.L., 599
Rust, R.W., 668
Ryan, J.D., 757, 1007

Saad, A.D., 206
Sailer, R.I., 140
Salom, S.M., 956

Sawyer, A.J., 575
Saxena, K.N., 967
Schaefer, P.W., 368
Schmidt, K.M., 471
Schowalter, T.D., 687, 984
Schroeder, W.J., 987
Schultz, P.B., 365
Schuster, D.J., 1001
Schwalbe, C.P., 202
Scott, D.R., 206, 535, 821
Sewall, D.K., 507
Shanks, C.H. Jr., 764
Shaver, T.N., 605
Shaw, P.B., 814
Shelton, A.M., 680
Shepard, B.M., 259
Shipp, J.L., 129, 683
Showers, W.B., 231, 779
Sinha, R.N., 618
Skinner, J.A., 922
Skoda, S.R., 902
Slessor, K.N., 905
Smith, C.M., 116
Smith, L.M., II, 971
Smith, S.M., 467
Sobhian, R., 645
Sorensen, E.L., 195, 708
Specht, H.B., 368
Spencer, N.R., 963
Stafford, K.C. III, 159
Stanek, E.J. III, 319
Starks, K.J., 676
Staten, R.T., 84
Stein, J.D., 563, 984
Stephen, F.M., 956
Sterling, W.L., 355
Stimac, J.L., 182
Stinner, B.R., 524
Stinner, R.E., 327
Stipanovic, R.D., 586
Stoffolano, J.G. Jr., 415
Stokes, M.E., 319
Stone, J.D., 877
Storey, G.K., 178
Story, J.M., 933
Story, R.N., 539
Stoyer, T.L., 864
Strickler, K., 30
Strnad, S.P., 975
Swift, F.C., 72
Sydnor, T.D., 996

Taylor, R.A.J., 1
Teng, P.S., 250
Terry, L.I., 237, 441, 625
Thoenes, S.C., 774
Thomas, G.D., 902
Thompson, C.R., 140
Thompson, D.C., 154
Thompson, L.C., 956
Thome, E.M., 731, 938
Thorarinsson, K., 593
Thorp, R.W., 664
Tipping, P.W., 809
Tollefson, J.J., 446
Torchio, P.F., 664
Torgersen, T.R., 90
Toscano, N.C. 25
Treacy, M.F., 420
Trosset, M.W., 791
Trumble, J.T., 374, 639
Tschinkel, W.R., 802
Turner, C.E., 111

Udevitz, M.S., 281
Underhill, E.W., 463
Unnithan, G.C., 967
Unruh, T.R., 979
Van Duyn, J.W., 237, 441, 625
Veal, R.A., 676
Villavaso, E.J., 951
Vinson, S.B., 747

Wagner, M.R., 551
Wagner, R.E., 639
Wagner, T.L., 390, 484, 497, 869, 951
Wallner, W.E., 556
Walmsley, M.H., 420
Walmsley, M.H., 471
Ward, R.H., 475
Waring, G.L., 15
Washburn, J.O., 286
Webster, J.A., 839
West, R.J., 905
Weston, P.A., 400
Whalon, M., 30
White, S.M., 535
Whitfield, G.H., 129, 683
Whitford, F., 231, 779
Wilkinson, R.C., 428
Williams, N.J., 747
Williams, R.E., 716
Williams, W.P., 532
Wilman, J.M., 990
Wilson, L.T., 545, 614, 1012
Wright, R.E., 211
Wu, H.-I, 497

Yaro, N.D., 126
Yearian, W.C., 691
Yendol, W.G., 459
Yonce, C.E., 68
Young, S.Y., 691

Zehnder, G.W., 135
Zoebisch, T.G., 1001